

Longleaf Pine Cone Prospects

for 2018 and 2019

Dale G. Brockway
Research Ecologist

Southern Research Station
USDA Forest Service
521 Devall Drive
Auburn, AL 36849

May 7, 2018

During the spring of 2018, cone production data were collected from selected low-density (e.g., shelterwood) stands of mature longleaf pine, throughout its native range. Binocular counts of green cones and unfertilized conelets were conducted on the crowns of sampled trees, as viewed from a single location on the ground. Visibility of cones and conelets on each tree is enhanced when the observer stands with their back to the sun. A breeze that moves the flexible pine needles about also helps the relatively more rigid cones and conelets stand out for the observer. The near-term regional averages and individual site averages for these counts are reported in Table 1.

Table 1. Estimated Longleaf Pine Cone Production.

Cooperator	State and County	Estimated cones per tree from green cones for fall 2018	Estimated cones per tree from conelets for fall 2019
Kisatchie National Forest	Louisiana, Grant	19.2	64.1
T.R. Miller Woodlands	Alabama, Escambia	5.4	60.8
Blackwater River State Forest	Florida, Santa Rosa	9.2	102.6
Eglin Air Force Base	Florida, Okaloosa	0.4	72.2
Apalachicola National Forest	Florida, Leon	0.8	92.9
Jones Ecological Research Center	Georgia, Baker	1.5	148.3
Tall Timbers Research Station	Florida, Leon	13.1	139.9
Fort Benning Military Base	Georgia, Chattahoochee	3.8	77.9
Sandhills State Forest	South Carolina, Chesterfield	13.9	83.7
Bladen Lakes State Forest	North Carolina, Bladen	12.0	89.1
Ordway-Swisher Biological Station	Florida, Putnam	0.4	19.9
Region Averages		7.2	86.5

Regional Summary:

The regional cone crop, based on green cone counts, is **failed for 2018**, at 7.2 cones per tree. The natural variation, typically seen throughout the native range of longleaf pine, is less evident in this year's data, with most sites failing and several other sites having only poor production. Bumper crops, good crops and fair crops were not observed anywhere. Poor crops (10 to 25 cones per tree) were present in Grant Parish, Louisiana, Leon County Florida, Chesterfield County South Carolina and Bladen County, North Carolina. Failed crops (< 10 cones per tree) were noted in Escambia County, Alabama, Santa Rosa County, Florida, Okaloosa County, Florida, Baker County, Georgia, Leon County, Florida, Chattahoochee County, Georgia and Putnam County, Florida.

The regional cone crop outlook, based on counts of unfertilized conelets, is **good for 2019**, at 86.5 cones per tree. The cone crop is forecasted to be a bumper crop at three sites, a good crop at seven sites and a poor crop at one site, reflecting a good deal of natural variability. However, keep in mind that cone crop estimates based on counts of unfertilized conelets are less reliable than those based on counts of green cones, because of conelet losses during their first year, with often fewer than half surviving to become green cones during their second year.

The 53-year regional cone production average for longleaf pine is about 28 green cones per tree. The single best cone crop occurred in 1996 and averaged 115 cones per tree. Good cone crops were observed in 1967 (65 cones per tree), 1973 (67 cones per tree), 1987 (65 cones per tree), 1993 (52 cones per tree), 2014 (98 cones per tree) and 2017 (62 cones per tree). Fair or better cone crops have occurred during 49% of all years since 1966, with an increased frequency since the mid-1980s. Reasons for this increasing frequency may be related to environmental change or management factors (or a combination of these). Research analysis of these long-term cone crop data has resulted in the recent publication of two scientific articles, which provide new insights into the reproductive pattern of longleaf pine in an environment with increasingly variable conditions. An electronic portable document file (pdf) of each of these two articles is included along with this report:

Guo, Q., Brockway, D.G., Chen, X., 2017. Temperature-related sex allocation shifts in a recovering keystone species, *Pinus palustris*. *Plant Ecology & Diversity* 10(4): 303-310.

Chen, X., Guo, Q., Brockway, D.G., 2017. Power laws in cone production of longleaf pine across its native range in the United States. *Sustainable Agriculture Research* 6(4): 64-73.

One additional article is currently in review and should be published in a scientific journal in the coming months:

Chen, X., Brockway, D.G., Guo, Q., 2018. Characterizing the dynamics of cone production for longleaf pine forests in the southeastern United States. *Forest Ecology and Management*. In Review.

Evaluating Longleaf Pine Cone Data:

Observations, concerning the natural variation in longleaf pine cone crops, and field studies, determining of the amount of seed (i.e., number of productive cones per tree) required to successfully regenerate even-aged shelterwood stands, resulted in development of Table 2. The minimum cone crop needed for successful natural regeneration, using an even-aged management technique such as the uniform shelterwood method, is 750 green cones per acre. This assumes 30 cones per tree, with 25 seed-bearing trees per acre. Thus, cone crops classified as “fair or better” represent regeneration opportunities, for which a receptive seedbed may be prepared through application of prescribed fire during the months prior to seed fall in October.

Table 2. Classification of Longleaf Pine Cone Crops*.

Crop Quality	Cones per Tree	Cones per Acre (on 25 trees per acre)
Bumper crop	≥ 100	≥ 2500
Good crop	50 to 99	1250 to 2475
Fair crop	25 to 49	625 to 1225
Poor crop	10 to 24	250 to 600
Failed crop	< 10	< 250

* Cones on mature trees (14-16 inches at dbh) in low-density stands (basal area < 40 feet²/acre).

When uneven-aged management stand-reproduction methods such as single-tree selection and group selection are being used, then “seed rain” incident on a site every year, although of variable intensity from year to year, is often sufficient for successful natural regeneration. While using selection silviculture frees one from dependency on the timing of good cone crops, it may nonetheless be useful for the manager of uneven-aged stands to be aware of cone crop quality from year to year when making management decisions.

It is also worth noting that a good deal of spatial variation occurs among longleaf pine stands across the Southern Region, relative to cone production. Therefore, even during a year with a lower overall regional average number of cones per tree, certain localities can experience substantial longleaf pine cone production. This regional report is intended as a guide, which broadly forecasts the overall status of longleaf pine cone production. Thus, we encourage forest managers to take binoculars to the field and carefully examine any individual stands in which they have an interest. In this way, they can, for those specific stands, acquire more detailed site-specific information that will aid them in making management decisions.

Study Partners:

Eddie Taylor, Kisatchie National Forest, Pineville, Louisiana
 Paul Padgett, T.R. Miller Woodlands, Brewton, Alabama
 Eric Howell, Blackwater River State Forest, Milton, Florida
 Alexander Sutsko, Natural Resources Management, Eglin Air Force Base, Niceville, Florida
 Ace Haddock, National Forests in Florida, Tallahassee, Florida
 Steve Jack, J.W. Jones Ecological Research Center, Newton, Georgia
 Eric Staller, Tall Timbers Research Station, Tallahassee, Florida
 James Parker, Land Management Branch, Fort Benning Military Base, Columbus, Georgia
 Brian Davis, Sandhills State Forest, Patrick, South Carolina
 Hans Rohr, Bladen Lakes State Forest, Elizabethtown, North Carolina
 Stephen Coates, Ordway-Swisher Biological Station, Melrose, Florida

Data Collection Cooperators:

Stephen Hudson, Land Management Branch, Fort Benning Military Base, Columbus, Georgia
Michael Low, Natural Resources Management, Eglin Air Force Base, Niceville, Florida
Lisa Huey, Ordway-Swisher Biological Station, University of Florida, Melrose, Florida
Alan Springer, Southern Research Station, USDA Forest Service, Pineville, Louisiana
Jacob Floyd, Southern Research Station, USDA Forest Service, Pineville, Louisiana
Cory Tucker, Southern Research Station, USDA Forest Service, Auburn, Alabama
Edward Andrews, Southern Research Station, USDA Forest Service, Auburn, Alabama

Cone Counting Method:

We have received many inquiries about the field method used when counting cones. Therefore, the following protocol and field data sheet are provided, in the event that you may wish to conduct your own observations of pine cone production in your locale. Remember:

- **Conelets** indicate how much production *may* happen next year (see Figure 1).
- **Green cones** tell you how much production will happen this year (see Figure 2)
- Brown cones tell you how much production occurred last year.



Figure 1. Two **conelets** on a longleaf pine branch, on either side of a bud.



Figure 2. Two **green cones** on a longleaf pine branch, as they would appear in spring.

- Equipment: 8 to 10x binoculars, field data sheet, clipboard, pencil, d-tape, tree tags, aluminum nails, orange flagging, bark scraper and orange tree paint.
1. Locate a stand that is growing at a shelterwood density of less than 40 square feet per acre (25 to 35 square feet per acre is a typical range) and contains numerous trees of at least 10 inches at dbh. Better cone crops come from larger-diameter trees and poorer cone crops come from smaller-diameter trees. A key consideration is that high brush and/or trees cannot obscure the crowns of your sample trees, or your data collection will be impaired. The midstory must be relatively open, so you can see the entire crowns of sample trees.

2. Select at least 10 trees in the stand to serve as your representative sample for monitoring, by painting a ring around the tree at dbh or higher and a sequence number on each (use a color other than white to avoid confusion with the white rings often painted around trees having RCW nests). You may also attach an aluminum tag to the tree, but attach this high enough so that the tag number will not become obscured by black char from or, even worse, melted during periodic prescribed fires (this happens when tags are too low).
3. Using the field data sheet, enter the following data at the top: location, date and crew. Then, for each tree, enter the tree number and its dbh. Now, you are ready to count the brown cones, green cones and small conelets.
4. While standing near each tree, count the number of brown cones lying on the ground around the tree. The cones from the most recent year appear brown and fresher than the cones from earlier years which appear weathered and gray. Enter this number. Then, walk toward the sun away from the tree. The precise distance away from the tree is not crucial, but it should be far enough away to give your neck a comfortable angle while looking up, but not so far away that you cannot clearly see the cones with 8 to 10 power binoculars. With the sun at your back, you may need to adjust your position a bit to the left or to the right, so that you can view the entire tree crown without moving from your counting location.
5. Work from the least difficult to most difficult strobili to see. First, count the number of brown cones still hanging on the tree from last fall. I usually start at the lower left of the crown and work my way up to the top of the crown, then across the top of the crown to the right and then down the right side of the crown all the way to the bottom-most branches. This is a systematic approach that sweeps across the entire crown (left half, top, right half) and leads to consistently accurate counts. Once you have done this, enter the number of brown cones still hanging on the tree into the data sheet.
6. Next, repeat the same up-over-down sweep with your binoculars, counting all of the green cones that can be seen from the single spot on which you are standing. Because these newer cones are green, they are more difficult to see against the green pine foliage. It helps to count these green cones (and other structures) on a bright sunny day, when the light is good. It also helps if there is a light breeze blowing that moves the pine needles about, thereby revealing the more rigid cones. Once you have done this, enter the number of green cones into the data sheet. This is perhaps the most important count you will make, since these green cones contain the seed that will be shed during the upcoming October, and it is these data that will become the numbers upon which the cone crop forecast for the current year will be based (a forecast in which many land managers have a great interest). News of a good cone crop usually alerts forest managers to get busy during the summer, preparing seedbeds that will be receptive to capturing and deriving the most benefit from the upcoming seed shed. You will also note on the data sheet that the raw number you see in your green cone count needs to be multiplied by 2 at the end of the column. Bill Boyer's research, through many years, confirmed that this adjustment to the raw count needed to be performed to obtain an accurate estimate (the actual regression from his work approximated 1.98). In general terms, he explained this as

being needed, because the cone count is performed by looking at only one side of the tree, thus the raw count for green cones needs to be doubled.

7. Finally, repeat the same up-over-down sweep with our binoculars, counting the small conelets that can be seen from the single spot on which you are standing. They are small, so this will take more time to locate them. But, they are up there. These conelets were pollinated only one month earlier (during March), but will not become fertilized for almost another 11 months (until a pollen tube grows from the surface of the conelet deep into its ovary). These conelets are the basis for estimating what the cone crop might be during the following year. But, it is worth bearing in mind that conelet abortion happens in nature for a variety of natural reasons (e.g., genetics, disease, insects, adverse weather conditions). Thus, not all conelets will survive to maturity. In fact, the conelet mortality rate is typically more than 50 percent. So, this estimate for next year, based on conelets, is less reliable than the forecast for this year, based on green cones.

Regional Longleaf Pine Cone Study: Female Strobili Count Data - - Field Data

Location: _____ Date: _____ Crew: _____

Tree Number	DBH	Brown Cones on Ground	Brown Cones on Tree	All Brown Cones	Green Cones	Conelets
1						
2						
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						

Total Count =

--	--	--	--

Adjusted Count performed only for Green Cones (is the Total Count x 2) =

Number Per Tree =

--	--	--

last year this year next year

Regional and Local


Summary and Graphs

Year	Southern Region	LA-Kisatchie National Forest	AL-Escambia Exp. Forest	W FL-Blackwater River State Forest	W FL-Eglin Air Force Base	W FL-Apalachicola National Forest	SW GA-Jones Res. Center	Red Hills-Tall Timbers Res. Station	W GA-Fort Benning Military Base	SC-Sandhills State Forest	NC-Bladen Lakes State Forest	FL Pen.-Ordway-Swisher Biological Station
1958			63.00									
1959			9.00									
1960			19.00									
1961			43.00									
1962			8.00									
1963			1.00									
1964			12.00									
1965			4.00									
1966	1.01		1.02			0.60						
1967	65.06	26.35	53.35	13.75		18.65	2.65					
1968	7.19	5.80	34.38	2.50	0.20	9.85	0.40				0.20	
1969	10.06	10.05	15.75	2.45	0.60	5.15	0.75			9.20	1.85	
1970	11.65	13.55	2.21	1.65	0.90	1.00	7.50			7.05	0.90	
1971	16.80	4.75	21.60	29.20	4.05	14.35	1.50			10.15	2.73	
1972	26.75	8.25	5.41	0.90	3.50	0.20	0.40			50.95	25.55	
1973	67.44	55.55	28.34	14.40	10.60	27.15	7.15			92.00	8.80	
1974	7.67	1.86	24.70	3.00	1.55	9.60	0.30			6.71	0.30	
1975	23.23		15.73	17.50	10.61		5.00			67.30		
1976	7.94		3.90	1.50	1.70	22.90	1.60			16.05		
1977	26.42	47.35	19.80	9.85	1.10	89.70	1.10			25.50	16.94	
1978	2.89	4.95	4.67	0.80	0.25	2.65	1.00			8.50	0.28	
1979	7.81	10.55	11.33	5.50	4.40		3.05			18.40	1.42	
1980	18.31	67.30	3.03	0.50	0.55		2.25			36.20		
1981	11.10	13.60	6.56	1.15	0.95		0.85			43.50		
1982	4.83	0.65	13.05	3.20	8.10		1.70			2.30		
1983	30.03	94.20	14.58	11.75	22.85		11.00			25.80		
1984	37.18	133.75	19.15	12.27	5.86		1.45			50.60		
1985	7.01	3.75	13.28	8.50	6.05		1.20			9.30		
1986	28.22	60.25	31.34	19.20	28.32		19.40			10.80		
1987	65.22	89.00	104.22	58.70	18.05		11.22			110.15		
1988	8.75	24.75	6.50	8.24			1.20			3.05		
1989	6.87	26.56	0.17	2.07			0.74			4.80		
1990	38.75	46.31	43.86	35.53			50.32			17.75		
1991	43.50	46.96	23.78	33.74			1.21			117.50	37.80	
1992	35.51	4.76	1.02	8.26		76.60	0.21			152.40	5.31	
1993	52.27	16.15	128.06	89.79		5.70	91.23		15.60	70.95	0.67	
1994	27.49	118.06	14.81	9.68	20.10	11.07	24.89			3.70	17.62	
1995	40.97	42.69	7.64	10.85	10.05	17.89	66.11		10.40	51.00	152.06	
1996	115.02	75.88	157.24	206.39	87.75	190.83	123.67		34.90	48.20	110.33	
1997	16.95	11.25	1.40	8.19	6.70	38.56	16.90		52.70	7.20	9.67	
1998	17.35	55.62	38.50	27.06	11.25	1.20	3.92		16.10	1.07	1.40	
1999	39.55	25.06	9.74	12.95	15.55	3.80	112.50	43.70	21.70	52.20	98.27	
2000	39.32	8.50	59.36	30.47	15.80	22.00	106.08	58.80	22.40	8.07	61.73	
2001	18.26	60.25	57.36	8.80	8.35	9.80	2.30	14.20	17.60	2.93	1.00	
2002	17.52	4.50	2.23	3.72	7.85	2.20	6.91	63.30	12.80	40.00	31.73	
2003	41.85	34.25	103.40	69.44	31.80	13.80	89.09	42.60	8.40	7.33	18.40	


2004	31.62	67.75	8.41	24.90	43.56	37.90	88.91	32.80	2.40	4.53	5.00	
2005	39.52	28.94	44.17	23.00	57.05	36.10	117.09	26.80	21.24	37.36	3.47	
2006	46.34	19.00	18.41	4.10	16.85	14.00	129.18	56.80		49.93	108.80	
2007	4.73	15.06	0.96	0.00	0.78	2.80	5.80	2.00	15.36	0.71	3.87	
2008	25.13	24.25	57.13	38.60	30.16	38.40	8.55	30.60	16.20	7.00	0.40	
2009	41.05	58.00	40.50	31.60	14.26	6.00	65.09	20.20	81.40	55.29	38.13	
2010	7.77	6.25	3.30	4.00	3.74	0.80	1.64	2.60	39.80	5.57	10.00	
2011	48.09	31.25	73.20	141.20	65.10	32.80	66.20	7.00	38.12	18.43	7.60	
2012	4.46	5.75	7.24	1.00	0.60	1.80	2.36	12.14	2.24	8.14	3.33	
2013	4.22	4.68	11.30	2.60	1.81	0.80	0.91	1.33	12.68	3.86	2.27	
2014	97.81	222.80	159.81	149.00	74.90	7.00	134.36	13.56	138.48	54.10	24.10	
2015	12.43	17.33	18.60	16.80	2.76	21.40	6.50	14.70	32.00	1.10	4.30	1.20
2016	3.38	1.47	0.48	1.00	2.64	0.60	1.45	3.78	5.92	6.86	12.53	0.40
2017	61.94	27.47	102.19	154.00	34.93	35.60	148.55	28.22	113.08	7.14	1.20	29.00
2018	7.23	19.20	5.43	9.20	0.36	0.80	1.45	13.11	3.76	13.86	12.00	0.40
Mean	27.91	36.05	29.71	26.62	15.44	21.33	29.94	24.41	30.64	29.25	22.16	7.75
Year	Southern Region	LA-Kisatchie National Forest	AL-Escambia Exp. Forest	W FL-Blackwater River State Forest	W FL-Eglin Air Force Base	W FL-Apalachicola National Forest	SW GA-Jones Res. Center	Red Hills-Tall Timbers Res. Station	W GA-Fort Benning Military Base	SC-Sandhills State Forest	NC-Bladen Lakes State Forest	FL Pen.-Ordway-Swisher Biological Station

Data are the average number of cones per longleaf pine tree forecasted for the fall (late October) with estimates based on counts of green cones during the spring (April and May) of each year.


Longleaf Pine Cone Production in Southern Region (since 1966)




Longleaf Pine Cone Production in Louisiana at Kisatchie NF (since 1967)




Longleaf Pine Cone Production in Southern Alabama at Escambia EF (since 1958)




Longleaf Pine Cone Production in West Florida at Blackwater River SF (since 1967)




Longleaf Pine Cone Production in Western Florida at Eglin AFB (since 1968)




Longleaf Pine Cone Production in Western Florida at Apalachicola NF (since 1966)




**Longleaf Pine Cone Production in Southwestern Georgia (since 1967):
at Southlands Forest Research Center from 1967 to 1996
and Jones Ecological Research Center since 1997**




**Longleaf Pine Cone Production in the Red Hills (since 1999):
at Pebble Hill Plantation from 1999 to 2009
and Tall Timbers Research Station since 2010**




Longleaf Pine Cone Production in Western Georgia at Fort Benning (since 1993)



Longleaf Pine Cone Production in South Carolina at Sandhills SF (since 1969)



Longleaf Pine Cone Production in North Carolina at Bladen Lakes SF (since 1968)



Longleaf Pine Cone Production on Florida Peninsula at Ordway-Swisher Biological Station (since 2015)

